

Laboratory Evaluation of the Effect of Egyptian Native Plants Against Some Parasitic Vectors

Mısır'daki Yerli Bitkilerin Bazı Parazitik Vektörlere Karşı Olan Etkisinin Laboratuvarında Değerlendirilmesi

Eman Taher¹, Narmeen Mahmoud¹, Maha Mahmoud²

¹Clinic of Parasitology, Research Institute of Ophthalmology, Giza, Egypt

²Department of Zoonoses, Faculty of Veterinary Medicine, Cairo University, Cairo, Egypt

ABSTRACT

Objective: Four plant extracts possessing molluscicidal and insecticidal efficacy were evaluated under laboratory conditions versus *Biomphalaria alexandrina*, *Lymnea cailliaudi* snails, their egg masses and *Culex pipiens* larvae. These extracts included Grape seed, *Eucalyptus*, *Pomegranate*, *Verbesina* alcoholic extracts, as well as *Eucalyptus* oil.

Methods: Different mortalities in the exposed vectors were recorded due to the four plant extracts using different concentrations and exposure time.

Results: Total snail mortality LC₁₀₀ was (100 ppm/12-24h) for Grape seed, (200 ppm/18-24h) for *Eucalyptus*, (100 ppm/12-18h) for *Pomegranate*, (100-200 ppm/24h) for *Verbesina* alcoholic extracts and (100-200 ppm/12h) for *Eucalyptus* oil. However, only *Eucalyptus*, *Verbesina* alcoholic extracts and *Eucalyptus* oil revealed snail ovicidal effects. LC₁₀₀ was (100-200 ppm/24h), (100-200 ppm/24h) & (100-200 ppm/12-48h) respectively. Moreover, the same plant extracts were able to induce total *Culex pipiens* larvicidal mortality, LC₁₀₀ was (200 ppm/48h). However, Grape seed and *Pomegranate* alcoholic extracts did not induce either snail ovicidal or *Culex pipiens* larvicidal total mortalities. Activities of the studied plant extracts were considered using reference molluscicidal (Copper sulfate) and insecticidal (Temephos) substances.

Conclusion: Egyptian native plants continue to provide a wealth of potential sources for biologically active agents that may have a promising role in the production of safe, biodegradable eco-friendly and natural molluscicidal and insecticidal agents.

(*Türkiye Parazit Derg* 2012; 36: 160-5)

Key Words: Molluscicides, insecticides, grape seed, eucalyptus, pomegranate, *Verbesina*, *Biomphalaria alexandrina*, *Lymnea cailliaudi*, *Culex pipiens*

Received: 24.02.2012

Accepted: 10.05.2012

ÖZET

Amaç: Mollusidal ve insektisidal etki gösteren dört bitki ekstratı laboratuvar koşulları altında *Biomphalaria alexandrina*, *Lymnea cailliaudi* salyangoz türleri, bunların yumurtaları ve *Culex pipiens*'in larvalarına karşı olan etkinlikleri açısından değerlendirilmiştir. Bu ekstraktlar, üzüm tohumu, ökaliptus, nar ve *Verbesina*'nın alkol ekstraktları ve ökaliptus yağıdır.

Yöntemler: Farklı konsantrasyonlar ve farklı sürelerde bu bitki ekstraktlarına maruz bırakılan vektörlerde farklı ölüm oranları elde edilmiştir.

Bulgular: Salyangozlardaki toplam ölüm oranları (LC₁₀₀) üzüm tohumu için 100 ppm/12-24saat, ökaliptus için 200 ppm/18-24saat, nar için 100 ppm/12-18saat, *Verbesina* için 100-200 ppm/24saat ve ökaliptüs yağı için 100-200 ppm/12saat olarak belirlenmiştir. Ancak sadece ökaliptus (LC₁₀₀=100-200 ppm/24saat) ve *Verbesina* LC₁₀₀=100-200 ppm/24saat) alkol ekstraktları ile ökaliptus yağı (LC₁₀₀=100-200 ppm/12-48saat) salyangoz yumurtaları üzerine öldürücü etki göstermiştir. Bundan başka bazı bitki ekstraktları *C. pipiens* larvaları üzerine olan öldürücü etkiyi indüklemiştir LC₁₀₀=200 ppm/48saat). Bununla birlikte, üzüm tohumu ve nar alkol ekstraktları ne salyangoz yumurtaları üzerine ne de *C. pipi-*

Address for Correspondence / Yazışma Adresi: Dr. Eman Taher, Clinic of Parasitology, Research Institute of Ophthalmology, Giza, Egypt

E-mail: eman_taher@yahoo.com

doi:10.5152/tpd.2012.38

ens larvaları üzerine etkili olmamıştır. Çalışılan bitki ekstraktlarının aktiviteleri referans mollusid (bakır sülfat) ve insektisit (Temefos) maddelere benzer olarak değerlendirilmiştir.

Sonuç: Mısır'daki yerli bitkilerin biyolojik olarak aktif vektörler için zengin potansiyel kaynaklar olduğu, güvenli, biyolojik olarak yıkılabilir, çevreye dost ve doğal mollusid ve insektisidal ajanlar üretilebileceği görülmüştür. (Türkiye Parazitolojî Dergî 2012; 36: 160-5)

Anahtar Sözcükler: Mollusid, insektisit, üzüm tahomu, ökaliptus, nar, *Verbesina*, *Biomphalaria alexandrina*, *Lymnea cailliaudi*, *Culex pipiens*

Geliş Tarihi: 24.02.2012

Kabul Tarihi: 10.05.2012

INTRODUCTION

Phytomedicine has been used to treat parasitism. Moreover; many modern commercial medicines are derived from plants. However, scientific evidence on the anti-parasitic efficacy of most plant products are limited (1). The use of natural products of plant origin (botanical derivatives) is a recent alternative approach for mosquito and snail control. In spite their toxicity to pests and snails, they are readily biodegradable and usually lack toxicity to higher animals' which means that they are eco-friendly (2).

Culex pipiens mosquito, *Biomphalaria alexandrina* and *Lymnea cailliaudi* snails are major pests of medical importance having different roles in transmitting many diseases (3).

Mosquitoes can transmit dreadful and fatal diseases to more than 700 million people, such as malaria, dengue, yellow fever, and filariasis (4). In various areas of the world, mosquitoes have been found to become resistant to several of the biological and conventional chemical insecticides used for their control (5).

Schistosomiasis is an important disease in Egypt and other tropical countries (6). It is considered as the world's most widespread parasitic disease (7). More than 207 million people are infected worldwide (8). A sure way to tackle the problem of diseases transmitted through snails as intermediate hosts is to destroy the carrier snails and remove an essential link in the life-cycle by using molluscicide (9). This type of control appears feasible and cost effective especially in poor countries (10). However, synthetic organic molluscicides are toxic to non-target animals and may have long term effects on the aquatic environment (9).

In this respect, the present work is an attempt to evaluate molluscicidal and or insecticidal properties of *Grape seed extract*, *Pomegranate*, *Verbesina*, *Eucalyptus* alcoholic extracts and *Eucalyptus oil* for their effect on *B. alexandrina* (vector of *Schistosoma mansoni*), *L. cailliaudi* (vector of *Fasciola gigantica*) snails and their egg masses as well as on larvae of *Culex pipiens*.

METHODS

Plant material

Grape seeds the fruit of *Vitis vinifera*, *Eucalyptus*, *Pomegranate* the fruit of *Punica granatum L.*, and *Verbesina alternifolia*, were collected from different areas from Egypt, during 2010. Plants were washed with tap water and dried at room temperature. The air-dried powdered parts were exhaustively extracted with 90% ethanol by maceration. The total alcoholic extract was combined and evaporated under reduced pressure by vacuum distillation at a temperature not exceeding 40°C to yield a semi-solid residue. Essential oil extract was obtained by hydrodistillation for 2 hours using a Clevenger-type apparatus using the methods of

Cetin and Yanikoglu (11). Investigation of the prepared oils was carried out on an Agilent (USA) GC-MS system. The extracts and oil were concentrated and stored in dark glass tubes under refrigeration at (4°C) until evaluation (12).

Tested Snails & their Egg Masses

Biomphalaria alexandrina and *Lymnea cailliaudi* snails (5-6 mm) were collected from irrigation canals in Giza Governorate. Snails were identified according to Christensen and Frandson (13). They were screened for natural infection with any trematodes. Uninfected snails that didn't harbor any trematoda parthenata were maintained in the laboratory conditions in fresh de-chlorinated tap water fed with fresh lettuce leaves with daily cleaning for at least 7 days before using in the experiment. Snails were prevented from crawling out of the solution by means of a fine mesh placed above the water surface.

Snail aquaria were supported by cellophane sheet for egg deposition. Freshly laid, 3-5 days old *B. alexandrina* and *Lymnea cailliaudi* egg masses were collected. Number of sound active embryos and their stage of development were identified microscopically per each snail egg mass.

Tested Mosquito

The vector mosquito, *Culex pipiens*, was used as test organism. Mosquito eggs were obtained from The Medical Research Institute of Insects. Eggs were soaked in de-chlorinated and filtered tap water to develop into first instar larvae. The larvae reared after this in the same aquarium till development to the third larval stage.

Molluscicidal and Larvicidal bioassay

The bioassay was done according to WHO (14) guidelines with slight modifications. Trematoda free; medium sized; active *B. alexandrina*, *L. cailliaudi* snails and their egg masses as well as late third instar larvae of *C. pipiens*; 30 in number for each group; were picked up from the aquaria and allocated in three 250 mL jars for screening the efficacy of the tested plant extracts.

Initially, 2 mL of each plant distillate was dissolved in 100 mL distilled water containing 0.3% Tween 80 to make 1% stock solution. Immersion technique was adopted according to WHO (14). Efficacy of each tested plant was evaluated using series of upgraded concentrations (400, 200, 100, 50, 25 and 12.5 p.p.m) each for (3, 6, 12, 24, 36, and 48) hours exposure time. Each experiment set contained three replicates. Control group was run simultaneously for each plant extract with the used solvent material in tap de-chlorinated water. All experiments were conducted simultaneously in the laboratory at room temperature (26±2°C).

At the end of the exposure time, tested solution was removed; snails, egg masses and larvae were washed for five times using

de-chlorinated tap water, kept in a new aquarium for 24 hours post exposure. Mortality was checked using crushing technique (5% sodium hydroxide solution) as described by WHO (15). Dose of plant extract in part per million (p.p.m) that killed 50% of the exposed stages (LC_{50}) and the dose that killed all of the exposed stages (LC_{100}); were calculated according to Kovendan et al. (16) using the following equation.

$$\text{Percentage mortality} = \frac{\text{Number of dead snails (larvae)} \times 100}{\text{Number of exposed snails (larvae)}}$$

RESULTS

Molluscicidal and larvicidal activities of four plant extracts included *Grape seed*, *Eucalyptus*, *Pomegranate*, as well as *Verbesina*, were tested at upgraded concentrations ranging from (12.5) to (400) p.p.m detecting LC_{50} and LC_{100} against *Biomphalaria alexandrina*; *Lymnea cailliaudi* snails and their egg masses as well as *Culex pipiens* larvae.

These extracts included *Grape seed extract*, *Pomegranate* alcoholic extract, *Verbesina* alcoholic extract, *Eucalyptus* alcoholic extract as well as, *Eucalyptus* oil. Copper sulphate and Temephos were included as reference molluscicidal and insecticidal substances (17, 18).

The tested plant extracts and oil proved to have molluscicidal activity but some of them had snail ovicidal as well as larvicidal activity versus *C. pipiens* larvae. Generally; there was a direct relationship between the efficacy and increasing the concentration and exposure time.

Results of the present study cleared that *Grape seed* extract was able to kill 50% of the exposed snails (LC_{50}) at a concentration of (100 ppm/6-12 h). These concentrations didn't show killing ability versus snail egg masses. In the same time total mortalities of the exposed snails (LC_{100}) were achieved using the same previous concentration after increasing the exposure time to 12-24h.

This extract showed weak effect versus *C. pipiens* larvae, mortality didn't increase over 50% after increasing the concentration more than (200 ppm for 48h).

Eucalyptus alcoholic extract induced 50% mortalities in the exposed snails and its egg masses at (100 ppm for 12-24h). Total mortality of *L. cailliaudi* snails and their egg masses was achieved at concentration of (100 ppm) and at (200 ppm) for *B. alexandrina* snails and their egg masses after 24h exposure time.

C. pipiens larvae showed more resistance, as they were able to tolerate higher concentrations of this material more than snails as LC_{50} & LC_{100} of this material reached up to (200 ppm after 24-48h) exposure time respectively.

The present study recorded rapid effect by *Pomegranate* alcoholic extract versus *L. cailliaudi* and *B. alexandrina* snails as 50% mortality was noticed at (100 ppm) after 3-6 hours exposure. On the contrary 50% mortalities (LC_{50}) for *L. cailliaudi* and *B. alexandrina* egg masses was recorded at (100-200 ppm) after 24h exposure. However, it was 200 ppm for *C. pipiens* larvae after 48h exposure. In the same time *Pomegranate* alcoholic extract couldn't induce total mortalities neither for both snails' egg masses nor for *C. pipiens* larvae.

Total mortalities (LC_{100}) of *L. cailliaudi* and its egg masses was revealed after using 100 ppm after 24h exposure of *Verbesina* alcoholic extract, higher concentration (200 ppm) for the same exposure time was required to get complete mortality for *B. alexandrina* snail and their egg masses. While 200 ppm after 48h was required for achieving total mortality of *C. pipiens* larvae.

The present study determined 100 ppm/12h as LC_{100} for *Eucalyptus* oil versus *L. cailliaudi* snails and its egg masses. While 200ppm was required for the LC_{100} versus *B. alexandrina*, after 12h. The concentration of (200 ppm/48h) was considered as LC_{100} of the oil versus *B. alexandrina* egg masses as well as *C. pipiens*.

It is worth to be mentioned that the molluscicidal activity of the studied plant extracts was higher than the molluscicidal activity of the reference material (copper sulfate) were its LC_{50} was (5-10 ppm) after 6-12 h exposure time and its LC_{100} reached up to (10-20 ppm) after 24 h. Moreover, the study recorded LC_{50} and LC_{100} of the reference insecticide (Temephos) as (0.5) after 24-48 h exposure time.

No mortalities were recorded in the control non exposed snails in water or in the used alcoholic solvent in each case. Results are presented in the Table 1.

DISCUSSION

Different snails and mosquitoes' species are of the most important vectors transmitting diseases all over the world; that constitute health problems and remaining as leading causes of mortality (19). Development of molluscicidal and insecticidal substances of botanical origin may serve as suitable alternatives to synthetic ones (17). Although several plants from different families have been reported for their molluscicidal and insecticidal activities, only few botanicals have moved from the laboratory to field use (20). In Egypt, screening of local plants for molluscicidal & mosquito larvicidal activities had received increasing attention (21). This was also previously reported by (22, 23).

Grape seed, *Eucalyptus*, *Pomegranate*, *Verbesina* alcoholic extracts, as well as *Eucalyptus* oil were evaluated in this study. The obtained results (LC_{100}) of the tested insecticidal activity of *Eucalyptus* was closely related to previous studies reported the larvicidal activity of *Eucalyptus* essential oil and others (24). It was also found that, essential oils of medicinal plants showed different mosquitocidal activities (25). Other studies also recorded that *Eucalyptus* alcoholic extract had molluscicidal activity (26). However, the present study revealed that *Eucalyptus* efficacy was less than that of *Grape seed* and *Pomegranate* alcoholic extracts. Moreover, *C. pipiens* larvae resisted its effect. *Eucalyptus* possesses insect and limited vermin control properties together with repellent properties. *Eucalyptus* alkaloids exhibited chemosterilant effect in addition to larvicidal and growth inhibition properties (27).

However, *Verbesina* induced 100% mortality after 48h with higher concentration than that was recorded by other studies (28). Medicinal uses of *Verbesina alternifolia* appeared to be limited and not widely documented although it was reported that it has an anti-inflammatory action. The major phyto-constituents are terpenoids, flavonoids and aromatic compounds (29). Poisoning

Table 1. Molluscicidal & insecticidal effect of plant extracts

No.	Tested Material	LC ₅₀					LC ₁₀₀				
		<i>B. alexandrina</i>		<i>L. cailliaudi</i>		<i>C. pipiens</i>	<i>B. alexandrina</i>		<i>L. cailliaudi</i>		<i>C. pipiens</i>
		snails	Egg masses	snail	Egg masses	Larvae	snails	Egg masses	snails	Egg masses	Larvae
1.	Grape Seed Extract	100ppm/ 12h	-ve	100ppm/ 6h	-ve	200ppm/ 48h	100ppm/ 24h	-ve	100ppm/ 12h	-ve	-ve
2.	Eucalyptus Alcoholic Extract	100ppm/ 12h	100ppm/ 24h	100ppm/ 12h	100ppm/ 12h	200ppm/ 24h	200ppm/ 18h	200ppm/ 24h	100ppm/ 24h	100ppm/ 24h	200ppm/ 48h
3.	Pomegranate Alcoholic Extract	100ppm/ 6h	200ppm/ 24h	100ppm/ 3h	100ppm/ 24h	200ppm/ 48h	100ppm/ 18h	-ve	100ppm/ 12h	-ve	-ve
4.	Verbesina Alcoholic Extract	200ppm/ 12h	-ve	100ppm/ 12h	100ppm/ 24h	200ppm/ 12h	200ppm/ 24h	200ppm/ 24h	100ppm/ 24h	100ppm/ 24h	200ppm/ 48h
5.	Eucalyptus oil	100ppm/ 12h	200ppm/ 24h	50ppm/ 24h	50ppm/ 12h	200ppm/ 24h	200ppm/ 12h	200ppm/ 48h	100ppm/ 12h	100ppm/ 12h	200ppm/ 48h
6.	Reference Molluscicide (Cu Sulphate)	10ppm/ 12h	10ppm/ 8h	5ppm/ 6h	10ppm/ 6h	-ve	20ppm/ 24h	20ppm/ 24h	10ppm/ 24h	10ppm/ 24h	-ve
7.	Reference Insecticide	-	-	-	-	0.5ppm/ 24h	-	-	-	-	0.5 ppm/ 48h

* In the same time no mortalities was recorded in control non exposed snails in water or in the used alcoholic solvent in each case

caused by *Verbesina* may be corresponded to high levels of nitrates and galegine (30).

The present study revealed that *Grape seed* and *Pomegranate* alcoholic extracts had potent molluscicidal activity. They caused significant behavioral changes in *B. alexandrina* and *L. cailliaudi* with the most obvious sign of distress being muscular and spiral twisting of the body, followed by crawling on one another. The nature and rapid onset of these behavioral responses shows the extracts probably contains neurotoxins, which amongst other things, might be active at the neuromuscular system of the exposed animals. Some of these characters were previously described (31).

In the author's opinion and in agreement with previous studies, Grape seed extract proved molluscicidal effect could be accepted as it possesses a wide array of pharmacological and biochemical actions (32). This includes anti-inflammatory, anti-carcinogenic activities. Moreover, it inhibits enzymes involved in the formation of histamine which might be attributed to its rich content of Proanthocyanidins.

Phytochemistry and pharmacological activities of *Pomegranate* (*Punica granatum*) were previously described (33). Medicinal properties are; antioxidant, anticancer, anti-inflammatory, anti-bacterial/antimicrobial and anthelmintic (34). Moreover, it was reported that *Pomegranate* extract gave very promising results as an anti-*trichomoniasis vaginalis* (35). *Pomegranate* contains high levels of phytochemicals including polyphenols, sugars, fatty acids, aromatic compounds, amino acids, tocopherols, sterols, terpenoids and alkaloids (36).

Generally, mortality to the plant distillates increased as concentration and exposure time increased (37). This could be due to that uptake of active moiety is time dependant, leading to progressive increase of entrance of the volatile oil and its effect in the snail body (20).

Molluscicides affect snails through inhibition of its respiratory enzymes either by direct contact or affecting the metabolic activities of snails (38). They act on different enzymes mainly those of respiration and carbohydrate metabolism. Inhibition of acetylcholinestrase enzyme, increasing its concentration at the synapses, leading to paralysis and eventually death (21). The adverse effect of these plant extracts can be accepted through such a mode of action. This was proved by appearance of *L. cailliaudi* as more sensitive than *B. alexandrina* snails.

This might be attributed to several factors; as *L. cailliaudi* has a wide aperture and large flappy body that needs more oxygen for the vital processes. As the majority of fresh water snails took their oxygen requirements throughout cutaneous pseudo bronchial and via pulmonary respiration. This is accompanied by high oxygen consumption level in comparison with *B. alexandrina* which has small snail body easily contracted inside snail with narrow aperture that protect its soft part of the body which is easily affected by the surrounding solution in comparison with the lower part of the body (snail foot) (21).

Moreover, sensitivity of *L. cailliaudi* egg masses in comparison with that of *B. alexandrina* may be related to that *L. cailliaudi* egg is large in size with thin transparent ootheca in comparison with that of *B. alexandrina* which is small in size with thick ootheca (21).

It is worth to be mentioned that bioactivity of plant extracts and essential oils is affected by the plant species ,cultivation conditions, plant storage, plant preparation and methods of extraction (39).

No behavioral symptoms or death occurred in control groups, indicating that no factors other than plant moieties were responsible for the altered behavior and mortality.

The present study revealed that all of the tested plant extracts had molluscicidal effect as well as snail ovicidal and *C. pipiens* larvicidal effect of some of them with high LC₅₀ and LC₁₀₀ in comparison with reference molluscicide (Copper Sulphate) and insecticide (Temephos) substances. This might be explained as the identified characters of the examined plant extracts were evaluated (on the level of the present study) as crude non-selectively concentrated extracts while the used reference substances were more purified. So free using of commercially purified, extraction of botanical molluscicidal and insecticidal active component, is considered to be a new field of preparation of safe, rapidly biodegradable and eco-friendly alternative to chemical ones (22, 40).

Acknowledgement

The authors are indebted and grateful to Dr. M. M. Salama, Department of Pharmacognosy, Faculty of Pharmacy, Cairo University, for preparing and supplying the plant extracts and for her support and advices.

Conflict of Interest

No conflict of interest was declared by the authors.

REFERENCES

- Githiori JB, Athanasiadou S, Thamsborg SM. Use of plants in novel approach for control of gastrointestinal helminthes in livestock with emphasis on small ruminants. *Vet Parasitol* 2006; 139: 308-20. [\[CrossRef\]](#)
- Shaalán EA, Canyon D, Younes MW, Abdel-Wahab H, Mansour AH. Effects of sub-lethal concentrations of synthetic insecticides and *Callitris glaucophylla* extracts on the development of *Aedes aegypti*. *J Vector Ecol* 2005; 30: 295-8.
- WHO (World Health Organization). The control of Schistosomiasis. Tech. Rep. 1985. 728, WHO, Geneva, Switzerland.
- Cetin H, Yanikoglu AA. Study of the larvicidal activity of *Origanum* (Labiatae) species from southwest Turkey. *J Vector Ecol* 2006; 31: 118-22. [\[CrossRef\]](#)
- Paul A, Harrington LC, Zhang L, Scott JG. Insecticide resistance in *Culex pipiens* from New York. *J Am Mosq Control Assoc* 2005; 21: 305-9. [\[CrossRef\]](#)
- Davis A. Schistosomiasis. In GC Cook, Manson's Tropical Diseases, 20th Edn., WB Saunders Company td, London 1996. PP: 1413-56.
- King CH & Dangerfield-Cha M. The unacknowledged impact of chronic schistosomiasis. *Chronic Illness* 2008; 4: 65-79. [\[CrossRef\]](#)
- Pereira AR, McCue CF, Gerwick WH. Cyanolide A, a Glycosidic Macrolide with Potent Molluscicidal Activity from the Papua New Guinea Cyanobacterium *Lyngbya bouillonii*. *J Nat Prod* 2010; 26: 217-20. [\[CrossRef\]](#)
- WHO (World Health Organization): "The control of Schistosomiasis". Tech. Rep. 1993. 830, WHO, Geneva, Switzerland.
- Lahlou M. Composition and molluscicidal properties of essential oils of five Moroccan Pinaceae. *Pharm Biol* 2003; 41: 207-10. [\[CrossRef\]](#)
- Cetin H, Yanikoglu A. A study of the larvicidal activity of *Origanum* (Labiatae) species from southwest Turkey. *J Vector Ecol* 2006; 31: 118-122. [\[CrossRef\]](#)
- Araújo EC, Silveira ER, Lima MA, Neto MA, de Andrade IL, Lima MA, et al. Insecticidal activity and chemical composition of volatile oils from *Hyptis martiusii* Benth. *J Agric Food Chem* 2003; 51: 3760-2. [\[CrossRef\]](#)
- Christensen NO & Frandsen A. An introduction to the taxonomy, morphology, biology and transmission ecology of species of the genus *Schistosoma* causing human African schistosomiasis. 1985. Danish Bilharsiasis Laboratory, Denmark.
- WHO Report of the WHO in formal consultation on the evaluation and on the testing of insecticides. 1996; CTD/WHO PES / IC/96.1, 69.
- World Health Organization Guidelines for laboratory and Field Testing of mosquito larvicides 2005; WHO.WHO/CDS/WHOPES/GCDPP.13WHO, Geneva, Switzerland.1-39.
- Kovendan K, Murugan K, Panneerselvam C, Kumar PM, Amerasan D, Subramaniam J, et al. Laboratory and field evaluation of medicinal plant extracts against filarial vector, *Culex quinquefasciatus* Say (Diptera: Culicidae). *Parasitol Res* 2011.
- Santos EA, de Carvalho CM, Costa ALS, Conceição AS, Moura FBP, Santana AEG. Bioactivity Evaluation of Plant Extracts Used in Indigenous Medicine against the Snail, *Biomphalaria glabrata*, and the Larvae of *Aedes aegypti*. *Evid Based Complement Alternat Med* 2012; 2012: 846583
- Subra R. Biology and control of *Culex pipiens* Quinquifasciatus Say 1823 (Diptera, Culicidae) with special reference to Africa. *Insect Sci Application* 1981; 1: 319-38.
- Kasi S, Komagata O, Tomita T, Sawabe K, Tsuda Y, Kurahashi H, et al. PCR-Identification of *Culex pipiens* complex collected in Japan. *Jpn J Infect Dis* 2008; 61: 184-91.
- Yadav RP, Singh A. Efficacy of *Euphorbia hirta* latex as plant derived molluscicides against freshwater snails. *Rev Inst Med Trop* 2011; 53: 101-6.
- Bakry FA. Use of Some Plant Extracts to Control *Biomphalaria alexandrina* Snails with Emphasis on Some Biological Effects. *World Appl Sci J* 2009; 6: 1335-45.
- Youssif RS, Shaalan EA. Mosquitocidal activity of some volatile oils against *Aedes caspius* mosquitoes. *J Vector Borne Dis* 2011; 48: 113-5.
- Oparaocha ET, Iwu I, Ahanakuc JE. Preliminary study on mosquito repellent and mosquitocidal activities of *Ocimum gratissimum* (L.) grown in eastern Nigeria. *Journal of Vector Borne Diseases* 2010; 47: 45-50.
- Traboulsi AF, EL-Haj S, Tueni M, Taoubi K, Nader NB, Mrad A. Repellency and toxicity of aromatic plant extracts against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). *Pest Manag Sci* 2005; 61: 596-604. [\[CrossRef\]](#)
- Prajapati V, Tripathi AK, Aggarwal KK, Khanuja SPS. Insecticidal, repellent and oviposition-deterrent activity of selected essential oils against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*. *Bioresour. Technol* 2005; 96: 1749-57. [\[CrossRef\]](#)
- Hammond JA, Fielding D, Nuru H. Eucalyptus: a sustainable self-delivery molluscicide? *Vet Res Commun* 1994; 18: 359-65. [\[CrossRef\]](#)
- Song A, Wang Y, Liu Y. Study on the chemical constituents of the essential oil of the leaves of *Eucalyptus globulus* Labill from China. *Asian Journal of Traditional Medicines* 2009; 4: 134-40.
- Amer A, Mehlhorn H. Larvicidal effect of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera, Culicidae). *Parasitol Res* 2006; 99: 466-72. [\[CrossRef\]](#)
- Sindhu, RK, Vasudeva N, Sharma SK. Pharmacognostical and Preliminary Phytochemical Investigations on *Verbesina encelioides* Benth. Roots. *Journal of Herbal Medicine and Toxicology* 2010; 4: 113-8.
- Shluker A. HNIS Report for *Verbesina encelioides* 2002.
- Singh A, Agarwal RA. Possibility of using latex of euphorbiales for snail control. *The Sci Total Environ* 1988; 77: 231-6. [\[CrossRef\]](#)
- Bagchi D, Sen CK, Ray SD, Das DK, Bagchi M, Preuss HG, Vinson JA. Molecular mechanisms of cardioprotection by a novel grape seed proanthocyanidin extract. *Mutat Res* 2003; 523-524: 87-97. [\[CrossRef\]](#)
- Lansky EP, Newman RA. *Punica granatum* (pomegranate) and its potential for prevention and treatment of inflammation and cancer. *Journal of Ethnopharmacology* 2007; 109: 177-206. [\[CrossRef\]](#)

34. Naovi SAH, Khan MSY, Vohora SB. Antibacterial, anti-fungal and anthelmintic investigations on Indian medicinal plants, *Fitoterapia* 1991; 62: 221.
35. El-Sherbini GM, Ibrahim KM, El Sherbiny ET, Abdel-Hady NM, Morsy TA. Efficacy of *Punica granatum* extract on in-vitro and in-vivo control of *Trichomonas vaginalis*. *J Egypt Soc Parasitol* 2010; 40: 229-44.
36. Chaturvedula Venkata, Sai Prakash & Indra Prakash Bioactive Chemical Constituents from Pomegranate (*Punica granatum*) Juice, Seed and Peel-A Review *Int J Res Chem Environ* 2011; 1: 1-18.
37. Cetin H, Yanikoglu A, Cilek JA. Larvicidal activity of selected plant hydrodistillate extracts against the house mosquito, *Culex pipiens*, a West Nile virus vector. *Parasitol Res* 2011; 108: 943-8. [\[CrossRef\]](#)
38. Singh SK, Singh A. Effects of the plants *Thevetia peruviana* and *Alstonia scholaris* (Family: Apocynaceae) on acetylcholinestrace activity of *Lymnea acuminata*. Egypt. *Egyptian J Schisto Infect End Dis* 2003; 25: 31-40.
39. Tawatsin AA, Thavara P, Wongsinkongman U, Bansidhi P, Boonruad J, Chavalit-tumrong T, et al. Repellency of essential oils extracted from plants in Thailand against four mosquito vectors (Diptera: Culicidae) and oviposition deterrent effects against *Aedes aegypti* (Diptera: Culicidae). *Southeast Asian J Trop Med Public Health* 2006; 37: 915-31.
40. Schall VT, Vasconcellos MC, Rocha RS, Souza CP, Mendes NM. The control of the schistosome-transmitting snail *Biomphalaria glabrata* by the plant Molluscicide *Euphorbia splendens* var. *hislopii* (syn *millii* Des. Moul): a longitudinal field study in an endemic area in Brazil. *Acta Trop* 2001; 79: 165-70. [\[CrossRef\]](#)